

# The effects of four forestry treatments on the community structure of spiders

Ferenc Samu<sup>1</sup>, Péter Ódor<sup>2</sup> and Zoltán Elek<sup>3</sup>

<sup>1</sup>MTA Centre for Agricultural Research, Agricultural Institute, Budapest, Hungary

<sup>2</sup>MTA Centre for Ecological Research, Institute of Ecology and Botany, Tihany, Hungary

<sup>3</sup>MTA-ELTE-MTM Ecology Research Group, Budapest, Hungary

## Motivation

Necessity of the harmonization between timber production and conservation purposes

Forest cover in Hungary: ~ 20%

Native: 63%

No management: 4%

Protected (management restrictions ): 24%

Silvicultural systems




- Shelterwood system (natural regeneration)
- Clear cutting system (artificial regeneration)
- Tree or group selection (new!, ~4%)





## Experimental design

- 70 yr old *Quercus petraea*, *Carpinus betulus* stand
- 5 treatments
  - thinning (d=80 m)
  - gap-felling (d=20 m)
  - clear-cutting (d=80 m)
  - retention-tree group (d=20 m)
  - control
- 6 replicates
- complete block design
- BACI (Before-After-Control-Impact)  
all measurements started in 2014


Pilis Mountains, N Hungary



## Spider density and species richness



## Community composition change



## Discussion

- Before treatment and two years after treatment states compared (2014-2016)
- Treatments created a gradient in light and humidity environment and in undergrowth cover
- All treatments resulted in a denser vegetation by 2016 compared to control
- Spiders responded to treatments by increased density and richness
- Community composition started to separate by treatments with greatest difference between control and clear-cut

## Acknowledgements

The project is supported by NKFI grant (OTKA) 111887  
We are grateful for assistance by Ms. Erika Botos

	2014	2015	2016
Adult abundance	1778	687	1654
Species number	21	21	22